Problem Set 0

Fall 11

Due: Tuesday, September 6th, in class, before class begins.

Please <u>follow</u> the homework format guidelines posted on the class web page:

```
http://www.cs.illinois.edu/class/fa11/cs373/
```

You must work on this problem set alone (groups are not allowed). Consult the following:

- Sipser Chapter 0.
- Chapters 1,2,3 of Discrete Mathematics by Chen (at http://rutherglen.science.mq.edu.au/wchen/lndmfolder/lndm.html)
- Lecture on Thursday, August 25th.
- 1. [Category: Notation, Points: 20]

Answer each of the following with "true", "false" or "wrong notation." Follow the notations in Sipser. $\{\ldots\}$ is used to represent sets and not multisets or anything else.

a) $\{a, b\} \cup \{a, b, c, d\} = \{a, b, c, d\}$ b) $\{a, b\} \cap \{c, d\} = \emptyset$ c) $\{a, b\} \cap \{c, d\} = \{\varnothing\}$ d) $\{a, b\} \setminus \{b, c\} = \{a, c\}$ e) $\{a, b\}/\{b, c\} = \{a\}$ $f) \ \emptyset \in \{a, b, c\}$ $g) \ \varnothing \subseteq \{a, b, c\}$ h) $\emptyset \in \emptyset$ i) $\varnothing \subset \varnothing$ j {} = {Ø} k) $\{a\} \subseteq \{\emptyset, a, b\}$ *l*) $\{a\} \subseteq \{a, \{a\}, \{\}\}$ $m) \ a \subseteq \{a, \{a\}, \{\}\}$ $n) \{a\} \in \{a, \{a\}, \{\}\}$ $o) \ a \in \{a, b, \{\}\}$ p) {a} × {b, c} = {(b, a), (c, a)} q) $\{a\} \times \{b, c\} = \{a, b\} \times \{c\}$ r) $|\{a, b\} \times \{c, d\}| = 4$

- s) $|powerset(\{a, b, c\})| = 3$
- t) $|\{a,b\}^3| = 3$
- 2. [Category: Relations, Points: 20]

Let $A = \{a, b, c\}$. Answer each of the following statements about relation on a set A with **true** or **false**. Explain your answer.

- a) The relation $R = \{(a, b), (b, a)\}$ is symmetric and reflexive.
- b) The relation $R = \{(a, a), (b, b), (c, c)\}$ is not symmetric and reflexive.
- c) The relation $R = \{(a, b), (b, c), (c, a)\}$ is symmetric and transitive.
- d) The relation $R = \{(a, b), (a, c), (a, a)\}$ is reflexive and transitive.
- e) The relation $R = \{(a, a), (b, b), (c, c)\}$ is symmetric, reflexive, and transitive.

Identify whether the following relations are equivalence relations. If not, state one property of equivalence relations that does not hold (reflexive, symmetric, transitive).

- (f) $R = \{(a, b) : a = b\}$ on the set \mathbb{N} .
- (g) $R = \{(a, b) : a < b\}$ on the set \mathbb{N} .
- (h) $R = \{(a, b) : a b \text{ is even}\}$ on the set \mathbb{N} .
- (i) $R = \{(a, b) : a + b \text{ is odd}\}$ on the set \mathbb{N} .
- (j) $R = \{(a, b) : a^2 = b^2\}$ on the set \mathbb{Z} .

3. [Category: Functions, Points: 10]

Determine whether each of the following functions is one-to-one or onto or both or neither. (Answers could be of the form "one-to-one but not onto" or "neither one-toone nor onto", etc.)

- a) $f_1: \mathbb{N} \to \mathbb{N}: x \mapsto x+1$
- b) $f_2: \mathbb{Z} \to \mathbb{Z}: x \mapsto 2x+1$
- c) $f_3: \mathbb{N} \to \mathbb{N}: x \mapsto x^2$
- d) $f_4: \mathbb{Z} \to \mathbb{Z}: x \mapsto x^2$
- e) $f_5: \mathbb{R} \to \mathbb{Z}: x \mapsto |x|$
- 4. [Category: Logic, Points: 20]

Let $\Sigma = \{a, b, \dots, z\}$. A finite sequence of symbols chosen from the alphabet Σ can be formally defined as a partial function $pos : \mathbb{N} \to \Sigma$, such that pos(n) is defined if and only if n does not exceed the length of the sequence. For example, the sequence abc can be defined as pos(1) = a, pos(2) = b, pos(3) = c, and pos(n) = undef for all n > 3.

If we write a first order logic (FOL) sentence over the above notations, it formally describes a property of finite sequences. For example, the sentence pos(5) = undef

says that the 5th position of the word is undefined, and hence the length of the sequence is less than 5. Also, the sentence " $\exists n \in \mathbb{N}(pos(n) = a \lor pos(n) = b \lor pos(n) = undef)$ " says that the sequence is entirely made up of a' and b's only.

Translate each of the following formal descriptions into an informal description as simple/natural as possible, e.g., using a short English sentence.

- a) $pos(3) \neq undef \land pos(4) = undef$
- b) $\forall n \in \mathbb{N} . pos(n) \neq a$
- c) $\exists n \in \mathbb{N} . pos(n) = a \land pos(n+1) = undef$
- d) $\not\exists n \in \mathbb{N} . pos(n) = a \land pos(n+1) = a$
- e) $\forall n \in \mathbb{N} . pos(n) = a \rightarrow pos(n+1) = b$

Translate each of the following English statements into a first-order logic sentence.

- (f) "the length of the sequence is at least 7."
- (g) "the sequence consists of purely a's."
- (h) "the sequence starts with 'ab'."
- (i) "every odd position in the sequence is an 'a'."
- (j) "the sequence contains a contiguous subsequence of the form 'a?c', where '?' indicates any single character."