

Optimization, Convexity, and Hyperplanes

We formulated the least squares method and linear regression as optimization problems. Our goal was to minimize the sum of the squared errors by choosing parameters for the linear model. Optimization problems have enormous utility in data science, and most model fitting techniques are cast as optimizations. In this chapter, we will develop a general framework for describing and solving several classes of optimization problems. We begin by reviewing the fundamentals of optimization. Next, we discuss convexity, a property that greatly simplifies the search for optimal solutions. Finally we derive vector expressions for common geometric constructs and show how linear systems give rise to convex problems.

5.1 Optimization

Optimization is the process of minimizing or maximizing a function by selecting values for a set of variables or parameters (called *decision variables*). If we are free to choose any values for the decision variables, the optimization problem is *unconstrained*. If our solutions must obey a set of constraints, the problem is a *constrained optimization*. In constrained optimization, any set of values for the decision variables that satisfies the constraints is called a *feasible solution*. The goal of constrained optimization is to select the “best” feasible solution.

Optimization problems are formulated as either minimizations or maximizations. We don’t need to discuss minimization and maximization separately, since minimizing $f(x)$ is equivalent to maximizing $-f(x)$. Any algorithm for minimizing can be used for maximizing by multiplying the objective by -1 , and vice versa. For the rest of this chapter, we’ll talk about minimizing functions. Keep in mind that everything we discuss can be applied to maximization problems by switching the sign of the objective.

During optimization we search for minima. A minimum can either be *locally* or *globally* minimal. A global minimum is has the




Figure 5.1: The maximum of a function $f(x)$ can be found by minimizing $-f(x)$.




Figure 5.2: Minima and maxima of a function can be local or global.

smallest objective value of any feasible solution. A local minimum has the smallest objective value for any of the feasible solutions in the surrounding area. The input to a function that yields the minimum is called the *argmin*, since it is the argument to the function that gives the minimum. Similarly, the *argmax* of a function is the input that gives the function's maximum. Consider the function $f(x) = 3 + (x - 2)^2$. This function has a single minimum, $f(2) = 3$. The minimum is 3, while the argmin is $x = 2$, the value of the decision variable at which the minimum occurs. For optimization problems, the minimum (or maximum) is called the *optimal objective value*. The argmin (or argmax) is called the *optimal solution*.

5.1.1 Unconstrained Optimization

You already know how to solve unconstrained optimization problems in a single variable: set the derivative to the function equal to zero and solve. This method of solution relies on the observation that both maxima and minima occur when the slope of a function is zero. However, it is important to remember that not all roots of the derivative are maxima or minima. Inflection points (where the derivative changes sign) also have derivatives equal to zero. You must always remember to test the root of the derivative to see if you've found a minimum, maximum, or inflection point. The easiest test involves the sign of the second derivative. If the second derivative at the point is positive, you've found a minimum. If it's negative, you've found a maximum. If the second derivative is zero, you've found an inflection point.

A similar approach works for optimizing multivariate functions. In this case one solves for points where the gradient is equal to zero, checking that you've not found an inflection point (called "saddle points" in higher dimensions).

5.1.2 Constrained Optimization

Constrained optimization problems cannot be solved by finding roots of the derivatives of the objective. Why? It is possible that the minima or maxima of the unconstrained problem lie outside the feasible region of the constrained problem. Consider our previous example of $f(x) = 3 + (x - 2)^2$, which we know has an argmin at $x = 2$. Say we want to solve the constrained problem

$$\min f(x) = 3 + (x - 2)^2 \quad \text{s.t.} \quad x \leq 1$$

The root of the derivative of f is still at $x = 2$, but values of x greater than one are not feasible. From the graph we can see that the minimum feasible value occurs at $x = 1$. The value of the derivative at $x = 1$ is -2 , not zero.




Figure 5.3: The function $f(x) = 3 + (x - 2)^2$ has a minimum of $f = 3$ at argmin $x = 2$.

Any point where the derivative of a function equals zero is called an *extreme point* or *extremum*. Setting the derivative of a function equal to zero and solving for the extrema is called *extremizing* a function.




Figure 5.4: The yellow region is the feasible space ($x \leq 1$). The global argmin occurs at $x = 1$. The derivative of the function is not zero at this point.

In general, constrained optimization is a challenging field. Finding global optima for constrained problems is an unsolved area of research, one which is beyond the scope of this course. However, there are classes of problems that we can solve to optimality using the tools of linear algebra. These problems form the basis of many advanced techniques in data science.

5.2 Convexity

Many “solvable” optimization problems rely on a property called *convexity*. Both sets and functions can be convex.

5.2.1 Convex sets

A set of points is *convex* if given any two points in the set, the line segment connecting these points lies entirely in the set. You can move from any point in the set to any other point in the set without leaving the set. Circles, spheres, and regular polygons are examples of convex sets.

To formally define convexity, we construct the line segment between any two points in the set.

Definition. A set S is convex if and only if given any $\mathbf{x} \in S$ and $\mathbf{y} \in S$ the points $\lambda\mathbf{x} + (1 - \lambda)\mathbf{y}$ are also in S for all scalars $\lambda \in [0, 1]$.

The expression $\lambda\mathbf{x} + (1 - \lambda)\mathbf{y}$ is called a *convex combination* of \mathbf{x} and \mathbf{y} . A convex combination of two points contains all points on the line segment between the two points. To see why, consider the 1-dimensional line segment between points 3 and 4.

$$\lambda(3) + (1 - \lambda)(4) = 4 - \lambda, \quad \lambda \in [0, 1]$$

When $\lambda = 0$, the value of the combination is 4. As λ moves from 0 to 1, the value of the combination moves from 4 to 3, covering all values in between.

Convex combinations work in higher dimensions as well. The convex combination of the vectors $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$ and $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$ is

$$\lambda \begin{pmatrix} 1 \\ 0 \end{pmatrix} + (1 - \lambda) \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} \lambda \\ 1 - \lambda \end{pmatrix}$$

The combination goes from the first point $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$ when $\lambda = 0$ to the second point $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$ when $\lambda = 1$. Halfway in between, $\lambda = 1/2$ and




Figure 5.5: The blue shapes are convex. The red shapes are not convex.




Figure 5.6: The segment connecting x and y can be defined as $\lambda x + (1 - \lambda)y$ for $\lambda \in [0, 1]$.

the combination is $\begin{pmatrix} 1/2 \\ 1/2 \end{pmatrix}$, which is midway along the line connecting $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$ and $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$. Sometimes it is helpful to think of a convex combination as a weighted sum of \mathbf{x} and \mathbf{y} . The weighting (provided by λ) moves the combination linearly from \mathbf{y} to \mathbf{x} as λ goes from 0 to 1.

5.2.2 Convex functions

There is a related definition for *convex functions*. This definition formalizes our visual idea of convexity (lines that curve upward) and concavity (lines that curve downward).

Definition 1. A function f is convex if and only if

$$f(\lambda\mathbf{x} + (1 - \lambda)\mathbf{y}) \leq \lambda f(\mathbf{x}) + (1 - \lambda)f(\mathbf{y}), \quad \lambda \in [0, 1]$$

This definition looks complicated, but the intuition is simple. If we plot a convex (upward curving) function, any chord – a segment drawn between two points on the line – should lie above the line. We can define the chord between any two points on the line, say $f(\mathbf{x})$ and $f(\mathbf{y})$ as a convex combination of these points, i.e. $\lambda f(\mathbf{x}) + (1 - \lambda)f(\mathbf{y})$. This is the right hand side of the above definition. For convex functions, we expect this chord to be greater than or equal to the function itself over the same interval. The interval is the segment from \mathbf{x} to \mathbf{y} , or the convex combination $\lambda\mathbf{x} + (1 - \lambda)\mathbf{y}$. The values of the function over this interval are therefore $f(\lambda\mathbf{x} + (1 - \lambda)\mathbf{y})$, which is the left hand side of the definition.

5.2.3 Convexity in Optimization

Why do we care about convexity? In general, finding local optima during optimization is easy; just pick a feasible point and move downward (during minimization) until you arrive at a local minimum. The truly hard part of optimization is finding global optima. How can you be assured that your local optimum is a global optimum unless you try out all points in the feasible space?

Fortunately, convexity solves the local vs. global challenge for many important problems, as we see with the following theorem.

Theorem. When minimizing a convex function over a convex set, all local minima are global minima.

Convex functions defined over convex sets must have a special shape where no *strictly* local minima exist. There can be multiple local minima, but all of these local minima must have the same value (which is the global minimum).




Figure 5.7: A convex combination in 2D.




Figure 5.8: Convex functions curve upward. Concave functions curve downward.




Figure 5.9: The chord connecting any two points of a convex function (red) lies above the function (blue).

All local minima are *less than or equal to* the global minimum. Strictly local minima must be *less than* the global minimum.

Let's prove that all local minima are global minima when minimizing a convex function over a convex set.

Proof. Suppose the convex function f has a local minimum at \mathbf{x}' that is not the global minimum (which is at \mathbf{x}^*). By the convexity of f ,

$$f(\lambda\mathbf{x}' + (1 - \lambda)\mathbf{x}^*) \leq \lambda f(\mathbf{x}') + (1 - \lambda)f(\mathbf{x}^*)$$

Since \mathbf{x}' is at a local, but not global, minimum, we know that $f(\mathbf{x}') > f(\mathbf{x}^*)$. If we replace $f(\mathbf{x}^*)$ on the right hand side by the larger quantity $f(\mathbf{x}')$, the inequality (\leq) becomes a strict inequality ($<$). (Even if both sides were equal, adding a small amount to the right hand side would still make it larger.) We now have

$$f(\lambda\mathbf{x}' + (1 - \lambda)\mathbf{x}^*) < \lambda f(\mathbf{x}') + (1 - \lambda)f(\mathbf{x}')$$

which, by simplifying the right hand side, becomes

$$f(\lambda\mathbf{x}' + (1 - \lambda)\mathbf{x}^*) < f(\mathbf{x}')$$

This statement says that the value of the function f on any point on the line segment from \mathbf{x}' to \mathbf{x}^* is less than the value of the function at \mathbf{x}' . If this is true, we can find a point arbitrarily close to \mathbf{x}' that is below our supposed local minimum $f(\mathbf{x}')$. Clearly, $f(\mathbf{x}')$ cannot be a local minimum if we can find a lower point arbitrarily closer to it. Our conclusion contradicts our original supposition. No local minimum can exist that are not equal to the global minimum. \square

For a simpler, yet less intuitive argument, let $\lambda = 1$. Then the inequality becomes $f(\mathbf{x}') < f(\mathbf{x}')$, which is nonsense.

The previous proof seemed to rely only on the convexity of the objective function, not on the convexity of the solution set. The role of convexity of the set is hidden. When we make an argument about a line drawn from the local to the global minimum, we assume that all the points on the line are feasible. Otherwise, it does not matter if they have a lower objective than the local minimum, since they would not be allowed. By assuming the solution set is convex, we are assured that any point on this line is also feasible.

5.2.4 Convexity of Linear Systems

This course focuses on linear functions and systems of linear equations. It would be enormously helpful if linear functions and the solution set of linear systems were convex. Then we can look for local optima during optimization and know that we've found global optima.

Let's first prove the convexity of linear functions. For a function to be convex, we require that a line segment connecting any two points in the line lie above or on the line. For linear functions, this is intuitively true. The line segment connecting any two points is the line

itself, so it always lies on the line. As a more formal argument, we describe a linear function as the product between a vector of coefficients \mathbf{c} and \mathbf{x} , i.e. $f(\mathbf{x}) = \mathbf{c}^T \mathbf{x}$. Let's start with the values of the function over the range spanned by arbitrary points \mathbf{x} and \mathbf{y} . The segment of the domain corresponds to the convex combination $\lambda \mathbf{x} + (1 - \lambda) \mathbf{y}$. The values of the function over this interval are

$$\begin{aligned} f(\lambda \mathbf{x} + (1 - \lambda) \mathbf{y}) &= \mathbf{c}^T (\lambda \mathbf{x} + (1 - \lambda) \mathbf{y}) \\ &= \mathbf{c}^T \lambda \mathbf{x} + \mathbf{c}^T (1 - \lambda) \mathbf{y} \\ &= \lambda \mathbf{c}^T \mathbf{x} + (1 - \lambda) \mathbf{c}^T \mathbf{y} \\ &= \lambda f(\mathbf{x}) + (1 - \lambda) f(\mathbf{y}) \end{aligned}$$

which satisfies the definition of convexity: $f(\lambda \mathbf{x} + (1 - \lambda) \mathbf{y}) \leq \lambda f(\mathbf{x}) + (1 - \lambda) f(\mathbf{y})$.

Now let's turn to a linear system $\mathbf{A}\mathbf{x} = \mathbf{b}$. We want to show that the set of all solutions for this system (the *solution space*) is convex. Let's assume we have two points in the solution space, \mathbf{x} and \mathbf{y} . Since \mathbf{x} and \mathbf{y} are solutions, we know that $\mathbf{A}\mathbf{x} = \mathbf{b}$ and $\mathbf{A}\mathbf{y} = \mathbf{b}$. If the solution set is convex, any point in the convex combination of \mathbf{x} and \mathbf{y} is also a solution.

$$\begin{aligned} \mathbf{A}(\lambda \mathbf{x} + (1 - \lambda) \mathbf{y}) &= \mathbf{A}\lambda \mathbf{x} + \mathbf{A}(1 - \lambda) \mathbf{y} \\ &= \lambda \mathbf{A}\mathbf{x} + (1 - \lambda) \mathbf{A}\mathbf{y} \\ &= \lambda \mathbf{b} + (1 - \lambda) \mathbf{b} \\ &= \mathbf{b} \end{aligned}$$

Since $\mathbf{A}(\lambda \mathbf{x} + (1 - \lambda) \mathbf{y}) = \mathbf{b}$, we know that all points on the line between \mathbf{x} and \mathbf{y} are solutions, so the solution set is convex.

5.3 Geometry of Linear Equations

Why do linear systems have convex solution spaces? Before answering, we should understand the shape of individual equations (rows) in the systems $\mathbf{A}\mathbf{x} = \mathbf{b}$. The equation corresponding to the i^{th} row is

$$\mathbf{A}(i, :) \cdot \mathbf{x} = b_i$$

which we will simplify by using the notation

$$\mathbf{a} \cdot \mathbf{x} = b$$

where \mathbf{a} and \mathbf{x} are vectors and the value b is a scalar. In two dimensions, this expression defines a line

$$a_1 x_1 + a_2 x_2 = b$$

By convention, all vectors are column vectors, including \mathbf{c} ; this requires a transposition to be conformable for multiplication by \mathbf{x} .

Following the conventions of the optimization field, we call the right hand side of linear systems the column vector \mathbf{b} , not \mathbf{y} as we have said previously.

The above representation of a line is the *standard form*, which differs from the *slope-intercept* form you remember from algebra ($y = mx + b$). It seems intuitive why the slope-intercept form is a line; a change in x produces a corresponding change $m\Delta x$ in y , with an intercept b when $x = 0$. What is the analogous reasoning for why $\mathbf{a} \cdot \mathbf{x} = b$ is a line?

First, we note that the vector \mathbf{a} always point perpendicular, or *normal* to the line. For the horizontal line $y = 3$, the vector $\mathbf{a} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$ points vertically. For the vertical line $x = 3$, the vector $\mathbf{a} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ point horizontal. For the line $x_1 + x_2 = 1$, $\mathbf{a} = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$, which is still perpendicular to the original line.

To help visualize the equation $\mathbf{a} \cdot \mathbf{x} = b$, we need to know the length of \mathbf{a} . The easiest solution is to normalize \mathbf{a} by dividing both sides of the equation by the norm of \mathbf{a} .

$$\frac{1}{\|\mathbf{a}\|} \mathbf{a} \cdot \mathbf{x} = b / \|\mathbf{a}\|$$

If we use our previous notation of $\hat{\mathbf{a}}$ for the normalized form of \mathbf{a} and define scalar $d = b / \|\mathbf{a}\|$, we have

$$\hat{\mathbf{a}} \cdot \mathbf{x} = d$$

We know that $\hat{\mathbf{a}}$ is a unit vector normal to the line. What is the meaning of d ? Let's compute the dot product $\hat{\mathbf{a}} \cdot \mathbf{x}$ using an arbitrary point \mathbf{x} on the line.

$$d = \hat{\mathbf{a}} \cdot \mathbf{x} = \|\hat{\mathbf{a}}\| \|\mathbf{x}\| \cos \theta = \|\mathbf{x}\| \cos \theta$$

Thus, d is the projection of the magnitude of \mathbf{x} onto the normal vector. For any point on the line, this projection is always the same length – the distance between the origin and the nearest point on the line. Conversely, a line is the set of all vectors whose projection against a vector $\hat{\mathbf{a}}$ is a constant distance (d) from the origin.

The same interpretation follows in higher dimensions. In 3D, the expression $\hat{\mathbf{a}} \cdot \mathbf{x} = d$ defines a plane with normal vector $\hat{\mathbf{a}}$ at a distance d from the origin. This definition fits with the algebraic definition of a plane that you may have seen previously: $a_1x_1 + a_2x_2 + a_3x_3 = d$. In higher dimensions (four or more), this construct is called a *hyperplane*.

Remember that when analyzing an expression of the form $\hat{\mathbf{a}} \cdot \mathbf{x} = d$, the constant on the right hand side (d) is only equal to the distance between the line and the origin if the vector $\hat{\mathbf{a}}$ is normalized. For example, the line

$$3x_1 + 4x_2 = 7$$




Figure 5.10: The vector \mathbf{a} is always normal (perpendicular) to the line $\mathbf{a} \cdot \mathbf{x} = b$.

The equation $\hat{\mathbf{a}} \cdot \mathbf{x} = d$ is called the Hesse normal form of a line, plane, or hyperplane.




Figure 5.11: A line is the set of all points \mathbf{x} whose projection onto $\hat{\mathbf{a}}$ is the distance d .

has coefficient vector $\mathbf{a} = \begin{pmatrix} 3 \\ 4 \end{pmatrix}$, which is not normalized. To normalize \mathbf{a} , we divide both sides by $\|\mathbf{a}\| = \sqrt{3^2 + 4^2} = 5$, yielding

$$\frac{3}{5}x_1 + \frac{4}{5}x_2 = \frac{7}{5}$$

Now we can say that the distance between this line and the origin is $7/5$.

5.4 Geometry of Linear Systems

The equation $\hat{\mathbf{a}} \cdot \mathbf{x} = d$ defines a hyperplane. It is also a single row in the linear system $\mathbf{Ax} = \mathbf{b}$. What does the entire system of equations look like? First, let's consider a set of three equations in two dimensions (so we can visualize them as lines). Solutions to $\mathbf{Ax} = \mathbf{b}$ are points of intersection of all three equations. If the lines are parallel, no solutions exist. If the lines all intersect at one point, there is a unique solution. If the lines are *colinear* (all fall upon the same line), infinitely many solutions exist. Note that these are the only options – zero, one, or infinitely many solutions, just as predicted by the grand solvability theorem. It is impossible to draw three straight lines that intersect in only two places.

If linear systems $\mathbf{Ax} = \mathbf{b}$ are a set of intersecting lines in 2D, what do the inequalities $\mathbf{Ax} \leq \mathbf{b}$ represent? Each inequality states that the projection of \mathbf{x} onto the normal vector \mathbf{a} must be less than d . These points form a *half-plane* – all the points on one side of a hyperplane. The system $\mathbf{Ax} \leq \mathbf{b}$ has a solution space that is the overlap of multiple half-planes (one for each row in \mathbf{A}). As we proved earlier, this solution set is a convex set.




Figure 5.12: A system of linear equations can have zero, one, or infinitely many points of intersection.




Figure 5.13: **A.** A single inequality defines a half-plane. **B.** Multiple half-planes intersect to form a convex solution set for the system $\mathbf{Ax} \leq \mathbf{b}$.