

Nonlinear Equations


How can we solve these equations?

- Spring force:

$$F = k x$$

What is the displacement when

$$F = 2\text{N}?$$




How can we solve these equations?

- Drag force:

$$F = 0.5 C_d \rho A v^2 = \mu_d v^2$$

What is the velocity when


$$F = 20\text{N?}$$



$$f(v) = \mu_d v^2 - F = 0$$



Find the root (zero) of the nonlinear equation $f(v)$




Nonlinear Equations in 1D


Goal: Solve $f(x) = 0$ for $f: \mathcal{R} \rightarrow \mathcal{R}$

Often called **Root Finding**

Bisection method



Bisection method



Convergence

An iterative method **converges with rate** r if:

$$\lim_{k \rightarrow \infty} \frac{\|e_{k+1}\|}{\|e_k\|^r} = C, \quad 0 < C < \infty \quad r = 1: \text{linear convergence}$$

Linear convergence gains a constant number of accurate digits each step
(and $C < 1$ matters!)

For example: Power Iteration

Convergence

An iterative method **converges with rate** r if:

$$\lim_{k \rightarrow \infty} \frac{\|e_{k+1}\|}{\|e_k\|^r} = C, \quad 0 < C < \infty$$

$r = 1$: linear convergence

$r > 1$: superlinear convergence

$r = 2$: quadratic convergence

Linear convergence gains a constant number of accurate digits each step
(and $C < 1$ matters!)

Quadratic convergence doubles the number of accurate digits in each step
(however it only starts making sense once $\|e_k\|$ is small (and C does not matter much))

Convergence

- The bisection method does not estimate x_k , the approximation of the desired root x . It instead finds an interval smaller than a given tolerance that contains the root.

Example:

Consider the nonlinear equation

$$f(x) = 0.5x^2 - 2$$


and solving $f(x) = 0$ using the Bisection Method. For each of the initial intervals below, how many iterations are required to ensure the root is accurate within 2^{-4} ?

A) $[-10, -1.8]$

B) $[-3, -2.1]$

C) $[-4, 1.9]$

Bisection method



Algorithm:

1. Take two points, a and b , on each side of the root such that $f(a)$ and $f(b)$ have opposite signs.
2. Calculate the midpoint $m = \frac{a+b}{2}$
3. Evaluate $f(m)$ and use m to replace either a or b , keeping the signs of the endpoints opposite.

Bisection Method - summary

- ❑ The function must be continuous with a root in the interval $[a, b]$
- ❑ Requires only one function evaluations for each iteration!
 - The first iteration requires two function evaluations.
- ❑ Given the initial internal $[a, b]$, the length of the interval after k iterations is $\frac{b-a}{2^k}$
- ❑ Has linear convergence


Newton's method

- Recall we want to solve $f(x) = 0$ for $f: \mathcal{R} \rightarrow \mathcal{R}$
- The Taylor expansion:

$$f(x_k + h) \approx f(x_k) + f'(x_k)h$$

gives a linear approximation for the nonlinear function f near x_k .

Newton's method



Example

Consider solving the nonlinear equation

$$5 = 2.0 e^x + x^2$$

What is the result of applying **one iteration** of Newton's method for solving nonlinear equations with initial starting guess $x_0 = 0$, i.e. what is x_1 ?

- A) -2
- B) 0.75
- C) -1.5
- D) 1.5
- E) 3.0

Newton's Method - summary

- ❑ Must be started with initial guess close enough to root (convergence is only local). Otherwise it may not converge at all.
- ❑ Requires function and first derivative evaluation at each iteration (think about two function evaluations)
- ❑ Typically has quadratic convergence


$$\lim_{k \rightarrow \infty} \frac{\|e_{k+1}\|}{\|e_k\|^2} = C, \quad 0 < C < \infty$$

- ❑ What can we do when the derivative evaluation is too costly (or difficult to evaluate)?

Secant method

Also derived from Taylor expansion, but instead of using $f'(x_k)$, it approximates the tangent with the secant line:

$$x_{k+1} = x_k - f(x_k)/f'(x_k)$$



Secant Method - summary

- ❑ Still local convergence
- ❑ Requires only one function evaluation per iteration (only the first iteration requires two function evaluations)
- ❑ Needs two starting guesses
- ❑ Has slower convergence than Newton's Method – superlinear convergence

$$\lim_{k \rightarrow \infty} \frac{\|e_{k+1}\|}{\|e_k\|^r} = C, \quad 1 < r < 2$$

1D methods for root finding:

Method	Update	Convergence	Cost
Bisection	Check signs of $f(a)$ and $f(b)$ $t_k = \frac{ b - a }{2^k}$	Linear ($r = 1$ and $c = 0.5$)	One function evaluation per iteration, no need to compute derivatives
Secant	$x_{k+1} = x_k + h$ $h = -f(x_k)/dfa$ $dfa = \frac{f(x_k) - f(x_{k-1})}{(x_k - x_{k-1})}$	Superlinear ($r = 1.618$), local convergence properties, convergence depends on the initial guess	One function evaluation per iteration (two evaluations for the initial guesses only), no need to compute derivatives
Newton	$x_{k+1} = x_k + h$ $h = -f(x_k)/f'(x_k)$	Quadratic ($r = 2$), local convergence properties, convergence depends on the initial guess	Two function evaluations per iteration, requires first order derivatives