

# Linear Programming II

## Lecture 19

October 29, 2015

## Simplex algorithm

```

Simplex(  $\hat{L}$  a LP )
  Transform  $\hat{L}$  into slack form.
  Let  $L$  be the resulting slack form.
   $L' \leftarrow \mathbf{Feasible}(L)$ 
   $x \leftarrow \mathbf{LPStartSolution}(L')$ 
   $x' \leftarrow \mathbf{SimplexInner}(L', x)$  (*)
   $z \leftarrow$  objective function value of  $x'$ 
  if  $z > 0$  then
    return "No solution"
   $x'' \leftarrow \mathbf{SimplexInner}(L, x')$ 
  return  $x''$ 
    
```

## Simplex algorithm...

1. **SimplexInner**: solves a LP if the trivial solution of assigning zero to all the nonbasic variables is feasible.
2.  $L' = \mathbf{Feasible}(L)$  returns a new LP with feasible solution.
3. Done by adding new variable  $x_0$  to each equality.
4. Set target function in  $L'$  to  $\mathbf{min} x_0$ .
5. original LP  $L$  feasible  $\iff$  LP  $L'$  has feasible solution with  $x_0 = 0$ .
6. Apply **SimplexInner** to  $L'$  and solution computed (for  $L'$ ) by **LPStartSolution**( $L'$ ).
7. If  $x_0 = 0$  then have a feasible solution to  $L$ .
8. Use solution in **SimplexInner** on  $L$ .
9. need to describe **SimplexInner**: solve LP in slack form given a feasible solution (all nonbasic vars assigned value 0).

## Notations

$B$  - Set of indices of basic variables  
 $N$  - Set of indices of nonbasic variables  
 $n = |N|$  - number of original variables  
 $b, c$  - two vectors of constants  
 $m = |B|$  - number of basic variables (i.e., number of inequalities)  
 $A = \{a_{ij}\}$  - The matrix of coefficients  
 $N \cup B = \{1, \dots, n + m\}$   
 $v$  - objective function constant.  
 LP in slack form is specified by a tuple  $(N, B, A, b, c, v)$ .

## The corresponding

$$\begin{aligned} \max \quad & z = v + \sum_{j \in N} c_j x_j, \\ \text{s.t.} \quad & x_i = b_i - \sum_{j \in N} a_{ij} x_j \text{ for } i \in B, \\ & x_i \geq 0, \quad \forall i = 1, \dots, n + m. \end{aligned}$$

5/26

## Reminder - basic/nonbasic

$$\begin{aligned} \max \quad & z = 29 - \frac{1}{9}x_3 - \frac{1}{9}x_5 - \frac{2}{9}x_6 \\ & x_1 = 8 + \frac{1}{6}x_3 + \frac{1}{6}x_5 - \frac{1}{3}x_6 \\ & x_2 = 4 - \frac{8}{3}x_3 - \frac{2}{3}x_5 + \frac{1}{3}x_6 \\ & x_4 = 18 - \frac{1}{2}x_3 + \frac{1}{2}x_5 \end{aligned}$$

Nonbasic variables

Basic variables

6/26

## The SimplexInner Algorithm

Description **SimplexInner** algorithm:

1. LP is in slack form.
2. Trivial solution  $x = \tau$  (i.e., all nonbasic variables zero), is feasible.
3. objective value for this solution is  $v$ .
4. Reminder: Objective function is  $z = v + \sum_{j \in N} c_j x_j$ .
5.  $x_e$ : nonbasic variable with positive coefficient in objective function.
6. Formally:  $e$  is one of the indices of  $\{j \mid c_j > 0, j \in N\}$ .
7.  $x_e$  is the **entering variable** (enters set of basic variables).
8. If increase value  $x_e$  (from current value of  $0$  in  $\tau$ )...
9. ... one of basic variables is going to vanish (i.e., become zero).

7/26

## Choosing the leaving variable

1.  $x_e$ : **entering variable**
2.  $x_l$ : **leaving** variable – vanishing basic variable.
3. increase value of  $x_e$  till  $x_l$  becomes zero.
4. How do we now which variable is  $x_l$ ?
5. set all nonbasic to  $0$  zero, except  $x_e$
6.  $x_i = b_i - a_{ie}x_e$ , for all  $i \in B$ .
7. Require:  $\forall i \in B \quad x_i = b_i - a_{ie}x_e \geq 0$ .
8.  $\implies x_e \leq (b_i/a_{ie})$
9.  $l = \arg \min_i b_i/a_{ie}$
10. If more than one achieves  $\min_i b_i/a_{ie}$ , just pick one.

8/26

## Pivoting on $x_e$ ...

1. Determined  $x_e$  and  $x_l$ .
2. Rewrite equation for  $x_l$  in LP.
  - 2.1 (Every basic variable has an equation in the LP!)
  - 2.2  $x_l = b_l - \sum_{j \in N} a_{lj} x_j$   
 $\implies x_e = \frac{b_l}{a_{le}} - \sum_{j \in N \cup \{l\}} \frac{a_{lj}}{a_{le}} x_j$ , where  $a_{ll} = 1$ .
3. Cleanup: remove all appearances (on right) in LP of  $x_e$ .
4. Substituting  $x_e$  into the other equalities, using above.
5. Alternatively, do Gaussian elimination remove any appearance of  $x_e$  on right side LP (including objective). Transfer  $x_l$  on the left side, to the right side.

9/26

## Pivoting continued...

1. End of this process: have new *equivalent LP*.
2. basic variables:  $B' = (B \setminus \{l\}) \cup \{e\}$
3. non-basic variables:  $N' = (N \setminus \{e\}) \cup \{l\}$ .
4. End of this **pivoting** stage:  
LP objective function value increased.
5. Made progress.
6. LP is completely defined by which variables are basic, and which are non-basic.
7. Pivoting never returns to a combination (of basic/non-basic variable) already visited.
8. ...because improve objective in each pivoting step.
9. Can do at most  $\binom{n+m}{n} \leq \left(\frac{n+m}{n} \cdot e\right)^n$ .
10. examples where  $2^n$  pivoting steps are needed.

10/26

## Simplex algorithm summary...

1. Each pivoting step takes polynomial time in  $n$  and  $m$ .
2. Running time of **Simplex** is exponential in the worst case.
3. In practice, **Simplex** is extremely fast.

11/26

## Degeneracies

1. **Simplex** might get stuck if one of the  $b_i$ s is zero.
2. More than  $> m$  hyperplanes (i.e., equalities) passes through the same point.
3. Result: might not be able to make any progress at all in a pivoting step.
4. Solution I: add tiny random noise to each coefficient.  
Can be done symbolically.  
Intuitively, the degeneracy, being a local phenomena on the polytope disappears with high probability.

12/26

## Degeneracies – cycling

1. Might get into cycling: a sequence of pivoting operations that do not improve the objective function, and the bases you get are cyclic (i.e., infinite loop).
2. Solution II: **Bland's rule**.  
Always choose the lowest index variable for entering and leaving out of the possible candidates.  
(Not prove why this work - but it does.)

13/26

## Correctness of

### Definition

A solution to an LP is a **basic solution** if it the result of setting all the nonbasic variables to zero.

**Simplex** algorithm deals only with basic solutions.

### Theorem

*For an arbitrary linear program, the following statements are true:*

- (A) *If there is no optimal solution, the problem is either infeasible or unbounded.*
- (B) *If a feasible solution exists, then a basic feasible solution exists.*
- (C) *If an optimal solution exists, then a basic optimal solution exists.*

Proof: is constructive by running the simplex algorithm.

14/26

## On the ellipsoid method and interior point methods

1. **Simplex** has exponential running time in the worst case.
2. **ellipsoid method** is *weakly* polynomial.  
It is polynomial in the number of bits of the input.
3. Khachian in 1979 came up with it. Useless in practice.
4. In 1984, Karmakar came up with a different method, called the *interior-point method*.
5. Also weakly polynomial. Quite useful in practice.
6. Result in arm race between the interior-point method and the simplex method.
7. BIG OPEN QUESTION: Is there *strongly* polynomial time algorithm for linear programming?

15/26

## Solving LPs without ever getting into a loop - symbolic perturbations

Details in the class notes.

16/26