

UNIVERSITY OF ILLINOIS, URBANA-CHAMPAIGN
Department of Electrical and Computer Engineering

ECE 313: Probability with Engineering Applications-Fall 2003

FIRST MIDTERM EXAM OCTOBER 15, 2003

Problem 1 (18 points) Let A and B be arbitrary events with $0 < P(A), P(B) < 1$. Indicate whether each of the following statements is true or false by clearly writing “true” or “false” on the line provided after each letter.

- (a) -----: $P(A \cup B) \leq P(A) + P(B)$. (d) -----: $P(A|B) + P(A^c|B) = 1$.
(b) -----: $P(A|B) \leq 1$. (e) -----: $\frac{P(AB)}{P(B)} \leq 1$.
(c) -----: $P(A|B) + P(A|B^c) = 1$. (f) -----: $P(AB^c) = P(A) - P(AB)$.

Problem 2 (12 points) X is a Binomial $(10, p)$ random variable. Compute the following:

- (a) The probability that X is equal to two. (b) The variance of $2 + 3X$.

Problem 3 (5 points) X is a random variable with mean and variance equal to unity: $\mu_X = \sigma_X^2 = 1$.

Compute $E[X^2]$.

Problem 4 (20 points) The random variables X and Y are independent, and each is uniformly distributed on the four integers $\{1, 2, 3, 4\}$. Compute the following:

- (a) $E[X]$, (b) $E[XY]$, (c) $P(X = k | XY = 4)$, $k = 1, 2, 3, 4$.

...Over

Problem 5 (20 points) A pair of dice is rolled until a sum of either 5 or 7 appears.

- (a) Let E_n denote the event that a 5 occurs on the n th roll and no 5 or 7 occurs on the first $n - 1$ rolls. Compute $P(E_n)$.
- (b) Compute the probability that a 5 occurs first.
- (c) Compute the probability that a 7 occurs first.

Problem 6 (25 points) A consumer electronics company purchases one million resistors from three different suppliers:

20% purchased from Supplier A, 30% from Supplier B, 50% from Supplier C

Of the resistors purchased, a certain number are defective, depending upon the supplier:

5% from Supplier A, 1% from Supplier B, 2% from Supplier C

A single resistor among these million is selected at random.

- (a) Compute $P(D | B)$ = the probability the resistor is defective, given it came from Supplier B .
- (b) Compute $P(D)$ = the probability the resistor is defective.
- (c) Compute $P(B | D)$ = the probability the resistor came from Supplier B , given that it is defective.
- (d) It is found that the resistor *is defective*. Express the maximum likelihood estimate of the supplier that sold this resistor, based on the quantities $P(D | B)$, $P(A | D)$, etc. You do not have to compute these conditional probabilities.
- (e) Express the MAP estimate of the supplier that sold the faulty resistor, in terms of the same quantities as in (d).

Useful fact: $1 + x + x^2 + \cdots = \sum_{k=0}^{\infty} x^k = \frac{1}{1-x}$ if $|x| < 1$.