Effective Lab Oral Report – Fall 2017

David Hertzog, Eugene V Colla, Virginia Lorenz
University of Illinois at Urbana-Champaign

We will present some of my slides and many Phys 403 student slides as examples. We can talk about why they are well constructed examples.

(All remarks about real slides are in these red boxes)

An eye-catching feature on slide 1

This is a technical presentation, so you must develop it as a logical sequence

Sentence title tells what the slide is about … the rest of the slide supports the assertion

- ☑ What was the goal?
  - What physics did you address?
  - What technology?
  - Define your special vocabulary here
- ☑ What did you actually do?
  - Apparatus / Procedures / Raw Data
- ☒ What are your results?
  - Polished graphs, proofs, numerical findings
  - Principal difficulties and uncertainties
- ☒ Conclusions

Fonts matter

Arial

Comic Sans

Times

Courier

Presentation components and grading scale.

- ✓ Title slide
- ✓ Science introduction
- ✓ Procedure
- ✓ Results. Analysis. Data.
- ✓ Conclusions. Suggestions etc.

CRITERIA		Score
Technical slides	(15)	
Science accuracy	(15)	
Quality of oral delivery and sharing of effort	(15)	
Got essential points across of effort	(15)	
Overall impression	(15)	
Final Totals	(75)	

Title slide

Title

OPTICAL STUDY OF FERROELECTRIC POTASSIUM DIDEUTERIUM PHOSPHATE (DKDP)



Author name

Student name

2/19/13

Date

Physics 403, Fall 2013
University of Illinois at Urbana-Champaign

Affiliation

Phase transition of Helium 4

- Below T_λ = 2.17 K, helium exists in mixture of superfluid and normal liquid helium.



The muon lifetime leads to the most precise determination of the Fermi constant, and gives the weak interaction strength

- The relation is

$$\frac{1}{\tau} \propto G_F^2 (1 + \delta)$$

measure this

- *MuLan* aims to determine τ_μ to 1 part per million precision, which requires:
  - 10^{12} muon decays
  - A muon beam of several MHz
  - A time-structured (chopped) beam



What happen if they are struck by pulses ?

A pulse or a series of pulses is used to change the net magnetization of system. Pulsed NMR!





90^0 Pulse



?

$$M(t) = M_0 e^{\frac{-t}{T_2}}$$

What happens to a nucleus in a magnetic field ?



(Courtesy of Bishop. K)

$$\Delta E = \gamma \cdot \hbar \cdot B_0 = \hbar\omega_0 \rightarrow \textit{Larmor frequency!}$$

Phase Transition in $BaTiO_3$









Phase Transition in $BaTiO_3$









Phase Transition in $BaTiO_3$









Phase Transition in $BaTiO_3$









ASIDE: Keep equations selective and informative

- What can an audience grasp in ‘real time’?
  - If they already know it, then they know it
  - If they don’t know it, they usually have to study it term by term
- Take a simpler approach
  - Substitute proportionalities for equalities ?
    - Can eliminates uninteresting constants
    - Can emphasize relationship of variables
  - Substitute words for blocks of standard terms?

$$\frac{1}{\tau} = \frac{G_F^2 m_\mu^5}{192\pi^3}(1+\delta)$$

$$\frac{1}{\tau} \propto G_F^2 (1+\delta)$$



Set them off attractively

$$\Gamma \propto (\text{phase space}) \times M_{ij}$$

  - Use builds and arrows to walk audience thru (see example)

Excitation and fluorescence signal convoluted together

observed signal

$$F(t) \propto \int_0^t E(t') F_\delta(t - t') dt'$$

excitation signal

simple fluorescence response (exponential)

- Excitation as sinusoid is simplest:

$$E(t) = E_0 + 2E_1 \cos(\omega t)$$

- Generalized through Fourier analysis
  - All periodic function can be expanded as sum of sinusoids

Show the equipment IF it helps explain your steps – not because you love it

- Photographs give scale and reality – but you add labels
- Schematics provide concept
- Icons strip away unnecessary details
- All of these techniques can be useful



Everybody loves an optical bench, but unless you map out the elements and the beam paths, it doesn't mean much

Experimental Apparatus



An example of image which is nice but does not help too much

Magnetic Field Calibration

- The magnetic field from the Earth and other residual magnetic fields is minimized by rotating the stand and adjusting the vertical field coils to minimize the zero field peak width.
- With the main field coils off, the sweep field is applied to determine the center of the zero field resonance (was found to be at 0.251A; using the geometry of the coils, this corresponds to 0.151 gauss).
- RF field is adjusted to provide maximum transition probability.



Too many words on slide

Setup diagrams, apparatus, measuring idea…



Schematic diagram adapted from notes

Samples: preparation, configuration etc.









Setup of Source and Detectors



Results

Raw tunneling data





Energy gap derived from tunneling conductivity

Count Rate vs. Pressure (Argon)

Data + fitting results α- range



Data + fitting results fluorescence



Difference in Up-Down (unnormalized)

Fit equation $Ne^{\frac{-t}{\tau}}\left(1+\alpha\cos(\omega t+\delta)\right)$



Courtesy Samuel Homiller and Pakpoom Buabthong Fall 2013

Difference in Up-Down (normalized)

Fit equation $Ne^{\frac{-t}{\tau}}\left(1+\alpha\cos(\omega t+\delta)\right)$



Courtesy Samuel Homiller and Pakpoom Buabthong Fall 2013

Results – witnessing a mystery?







Presenting data is your most important and challenging task







Fitting to the Curie-Weiss law



$$\varepsilon' = \frac{C}{T - T_C} + off$$

$C = 3563.3 \pm 0.4\,\mathrm{K}$

$T_C = 118.9825 \pm 0.0003\,\mathrm{K}$

Courtesy Zongyuan Wang and Arnulf Taylor Su 2017

AFM of Optical Data Storage Media

CD



DVD



Blu-Ray



	CD	DVD	Blu-Ray
Mark length	0.99 - 2.96	0.48 - 1.45	0.14 - 0.41
Track pitch	1.63	1.00	0.40
Track width	0.50	0.24	0.15

Units in µm

Fitting the data

$$V = C\sqrt{\left(\frac{T - T_{offset}}{T_\lambda}\right)\left(1 - \left(\frac{T - T_{offset}}{T_\lambda}\right)^{5.6}\right)} \quad \rightarrow \quad V = C\left[\left(\frac{T - T_{offset}}{T_\lambda}\right)\left(1 - \left(\frac{T - T_{offset}}{T_\lambda}\right)^{\beta}\right)\right]^{\gamma}$$

Offset, intrinsic to the experiment

Fit to the exponents as well

$$C \approx 26$$

$$T_\lambda \approx 2.17$$



Perform the 5 parameter fit-

The values that are obtained are not very close to the expected values

Also, the fit is not the best

Try to fit the data with this function

$$V = \left(1 - \frac{T - T_{offset}}{T_{\lambda}}\right)^{\gamma}$$

The data refuses to fit to this function



Finish your talk with the data analysis and conclusions and a slide showing the main points you want us to remember

- **Make sure you discuss the principal uncertainties.**
  - ***For most of these experiments, it will be how accurately does your instrument measure something***
  - ***A few experiments will also have statistical uncertainties … more data leading to a better finding***
- **Include a representative (simplified) graphic**
  - ***This slide will be up during question period so this graphic will get burned into people's memory***
- **Because this is a lab, offer some advice for others who follow**